

PACKAGE LEAFLET: INFORMATION FOR THE USER

Locoid Crelo 0.1% cutaneous emulsion

Hydrocortisone butyrate

Read all of this leaflet carefully before you start using this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please inform your doctor or pharmacist.

In this leaflet:

1. What Locoid Crelo is and what it is used for
2. Before you use Locoid Crelo
3. How to use Locoid Crelo
4. Possible adverse reactions
5. How to store Locoid Crelo
6. Further information

1. WHAT LOCOID CRELO IS AND WHAT IT IS USED FOR

Locoid Crelo contains a hormone of the adrenal cortex (corticosteroid) and quickly reduces the symptoms of some skin diseases – itching, redness, skin scaling. Corticosteroids do not treat the main disease.

Locoid Crelo may be used for treatment of superficial skin conditions (not caused by microorganisms), often accompanied with itching, redness or skin scaling. Locoid Crelo is especially suitable for treatment of hairy skin.

2. BEFORE YOU USE LOCOID CRELO

Do not use Locoid Crelo:

- If you have skin diseases, caused by infections, which derive from bacteria, viruses, fungi, yeast or parasites. This diseases may get complicated or be masked.
- If you have acne, as well as skin redness.
- If you have wounds or ichthyosis.

- If you have skin diseases which may be a result of previous treatment with corticosteroids as skin inflammation around mouth, skin thinning , possibly with striae, fragil blood vessels.
- In case of rash in children with sore of walking feet.
- If you allergic (hypersensitive) to hydrocortisone butyrate or some of the other Locoid Crelo ingredients (with occurs rarely). See also Section 'Important Information' regarding some Locoid Crelo ingredients.

Take special care with the use of Locoid Crelo

- Facial skin and genitals area skin, as well as in skin folders is especially sensitive to corticosteroids. That is why it should be treated with care.
- If you apply Locoid Crelo on a large skin area or in children, a physician examination is necessary.
- Locoid Crelo should not get into the eyes and should not by applied on eyelids.
- Continuous corticosteroids use on a large skin area should be avoided in children..

Consult your doctor if some of the above mentioned warnings concern you.

If after continuous treatment with Locoid Crelo the disease returns after a short time, you should consult your doctor.

Use of other drugs

Please, note: The following remarks may also refer to the use of drugs in the near past or near future. Please, inform your doctor or pharmacists if you take or have recently taken other drugs, including OTC drugs.

Pregnancy and lactation

Pregnancy:

The use of high doses of corticosteroids may lead to retardation of the foetus growth. As only a small quantity of the active substance is available in blood during the use of Locoid Crelo , it may be used also for a short time on small skin areas. The above described effect cannot be excluded in continuous treatment or in application on larger skin areas.

Ask your doctor or pharmacist for an advice before the use of any drug. If you want to use Locoid Crelo during pregnancy you should consult first your doctor.

Lactation:

Locoid Crelo may be used by women who breast-feed in case of short-term treatment and the product is applied on small areas. In continuous treatment or in treating of large skin areas breast-feeding is not recommended.

Ask your doctor or pharmacist for an advice before use of any drug. If you want to use Locoid Crelo during breast-feeding you should consult first your doctor.

Effects on ability to drive and use machines

There is no data for the effect of Locoid Crelo on the ability to drive and use machines, but it is not expected Locoid Crelo to have such an effect.

Important information regarding some Locoid Crelo ingredients

Cetostearyl alcohol may provoke local skin reactions (e.g. contact dermatitis). Propyl parahydroxibenzoate (E 216) and butyl parahydroxibenzoate (E 218) may provoke allergic reactions (possibly delayed). Butylhydroxitoluen (E321) may provoke local skin reactions (e.g. contact dermatitis) or eyes or mucosa irritation.

3. HOW TO USE LOCID CRELO

Locoid Crelo should be applied in a thin layer on the affected skin. After that it may be massaged carefully. It should not be applied more than 1 to 3 times daily on the affected skin, if your doctor has not prescribed other.

If complaints reduce, your doctor may advise you to use Locoid Crelo rarely. Usually you should not use more than 1-2 vials of 30 g Locoid Crelo a week.

If you have used more Locoid Crelo than you should

If you have used more than the necessary dose of Locoid Crelo, adverse reactions (as described in Section 4 'Possible adverse reactions') can occur. In this case contact your doctor or pharmacist immediately.

If you have missed to use Locoid Crelo

Once you have forgotten to apply Locoid Crelo, continue using Locoid Crelo as usual. No need to apply higher quantity of Locoid Crelo.

If you have stopped using Locoid Crelo

If Locoid Crelo treatment is stopped too early, primary symptoms may return. Always consult your doctor if you intend to stop treatment.

4. POSSIBLE ADVERSE REACTIONS

Like all medicines Locoid Crelo may cause adverse reactions, although not everybody gets them.

During Locoid Crelo application the following adverse reactions may occur:

Immune system disorders

Very rare (in less than 1 in 10 000 patients)

- hypersensitivity

Metabolic system disorders

Rare (in more than 1 in 10,000, but less than 1 in 1,000 patients)

- adreno-cortical suppression (suppression of the adrenal gland)

In case of continuous treatment with corticosteroids applied on the skin, they may penetrate through the skin and reach blood. This may lead to adreno-cortical suppression. There is a higher risk of systemic adverse reactions in case of:

- application under occlusive dressing (under occlusion, in skin folds)
- application on large skin areas
- continuous treatment
- application in children

Eyes disorders

Very rare (in less than 1 in 10 000 patients)

Increased intra-ocular pressure with increased risk of cataract.

Skin and subcutaneous tissue disorders

Rare (in more than 1 in 10,000, but less than 1 in 1,000 patients)

- burning or itching sensation of the treated with Locoid Crelo skin. Usually this is not serious and disappears by itself.
- Skin discoloration and thinning (often irreversible), sometimes with striae, skin blood vessels dilatation, excessive hairiness (hypertrichosis).
- Worsening of a skin disease after discontinuation of the treatment ('rebound effect').
- Itching, pimples or purulence if you have used continuously Locoid Crelo under occlusive dressing.
- Ekzema or skin inflammation (dermatitis), including skin inflammation as a result of a contact with irritating substances or substances provoking hypersensitivity (contact dermatitis).

If any of the adverse drug reactions gets serious, or if you notice any other reactions, not listed in this leaflet, please inform your doctor or pharmacist.

5. HOW TO STORE LOCOID CRELO

Keep out of the reach and sight of children

Locoid Crelo should be stored under 25° C

Locoid Crelo should not be used after the expiry date, mentioned on the vial and the carton package. The expiry date refers to the last day of the mentioned month.

Drugs should not be thrown in wastewater or in household waste container. Ask your pharmacist how to destroy the drugs you no longer use. These measures will help to protect the environment.

6. FURTHER INFORMATION

What does Locoid Crelo contain:

The active substance is hydrocortisone butyrate.

1 g Locoid Crelo cutaneous emulsion contains the active substance *hydrocortisone butyrate* 1 mg.

Other ingredients are: cetostearyl alcohol, macrogol cetostearylether, white soft paraffin, hard paraffin, borage, refined oil, butylhydroxytoluene E321, propyleneglycol, E1520, sodium citrate, anhydrous E331, citric acid, anhydrous E330, propyl parahydroxybenzoate E216, butyl parahydroxybenzoate E218, purified water.

How Locoid Crelo looks like and what the package contains

Locoid Crelo is white cutaneous emulsion. It is in polyethylene vials with polyethylene caps 30 g. The vials is packed in yellow/grey/white carton box.

Marketing Authorization Holder:

LEO Pharma A/S
Industriparken 55
DK-2750 Ballerup
Denmark

Manufacturers:

Astellas Pharma Europe BV
Sylviusweg 62
2333 BE Leiden
The Netherland

and

Temmler Italia S.R.L.
Via delle Industrie 2
20061 Carugate (MI)
Italy

Date of the last leaflet approval

07/2016